arXiv:1304.5447v1 [math.AG] 19 Apr 2013

ON A REPRESENTATION OF THE FUNDAMENTAL CLASS OF AN IDEAL DUE TO LEJEUNE-JALABERT

ELIZABETH WULCAN

Abstract. Lejeune-Jalabert showed that the fundamental class of a Cohen-Macaulay ideal $\mathfrak{a} \subset \mathcal{O}_0$ admits a representation as a residue, constructed from a free resolution of $\mathfrak{a}$, multiplied by a certain differential form coming from the resolution. We give an explicit description of this differential form in the case when the free resolution is the Scarf resolution of a generic monomial ideal. As a consequence we get a new proof and a refinement of Lejeune-Jalabert's result in this case.

1. Introduction

In [L-J] Lejeune-Jalabert showed that the fundamental class of a Cohen-Macaulay ideal $\mathfrak{a}$ in the ring of germs of holomorphic functions $\mathcal{O}_0$ at $0 \in \mathbf{C}^n$ admits a representation as a residue, constructed from a free resolution of $\mathfrak{a}$, multiplied by a certain differential form coming from the resolution, see also [L-J2, AL-J]. This representation generalizes the well-known fact that if $\mathfrak{a}$ is generated by a regular sequence $f = (f_1, \ldots, f_n)$, then

$$\operatorname{res}_f(df_n \wedge \cdots \wedge df_1) = \frac{1}{(2\pi i)^n} \int_\Gamma \frac{df_n \wedge \cdots \wedge df_1}{f_n \cdots f_1} = \dim_{\mathbf{C}}(\mathcal{O}_0/\mathfrak{a}), \tag{1.1}$$

where res_f is the *Grothendieck residue* of $f_1, \ldots, f_n$ and $\Gamma = \{|f_j| = \epsilon\}$ for some ϵ such that f_j are defined in a neighborhood of $\{|f_j| \leq \epsilon\}$, see [GH, Chapter 5.2].

We will present a formulation of Lejeune-Jalabert's result in terms of currents. Recall that the *fundamental cycle* of $\mathfrak{a}$ is the cycle

$$[\mathfrak{a}] = \sum m_j [Z_j],$$

where Z_j are the irreducible components of the variety Z of $\mathfrak{a}$, and m_j are the *geometric multiplicites* of Z_j in Z, defined as the length of the Artinian ring $\mathcal{O}_{Z_j,Z}$, see, e.g., [F, Chapter 1.5]. In particular, if $Z = \{0\}$, then $[\mathfrak{a}] = \dim_{\mathbf{C}}(\mathcal{O}_0/\mathfrak{a})[\{0\}]$.

Assume that

$$0 \to E_p \xrightarrow{\varphi_p} \ldots \xrightarrow{\varphi_2} E_1 \xrightarrow{\varphi_1} E_0 \tag{1.2}$$

is a free resolution of of $\mathcal{O}_0/\mathfrak{a}$ of minimal length $p := \operatorname{codim} \mathfrak{a}$; here the E_k are free $\mathcal{O}_0$-modules and $E_0 \cong \mathcal{O}_0$. In [AW], together with Andersson we constructed from (1.2) a (residue) current R, which has support on Z, takes values in E_p, and is of bidegree $(0,p)$. If (1.2) is the Koszul complex of a regular sequence $f = (f_1, \ldots, f_p)$, then R coincides with the *Coleff-Herrera residue current* $R^f_{CH} = \bar\partial[1/f_1] \wedge \cdots \wedge \bar\partial[1/f_p]$ of f, introduced in [CH]. Given bases of E_k, let $d\varphi_k$ be the $\operatorname{Hom}(E_k, E_{k-1})$-valued

---

Date: April 22, 2013.

The author is supported by the Swedish Research Council.

$(1,0)$-form with entries $(d\varphi_k)_{ij} = d(\varphi_k)_{ij}$ if $(\varphi_k)_{ij}$ are the entries of φ_k and let $d\varphi$ denote the E_p^*-valued $(p,0)$-form

$$d\varphi := d\varphi_1 \circ \cdots \circ d\varphi_p.$$

Now, identifying $[\mathfrak{a}]$ with the current of integration along $[\mathfrak{a}]$, see, e.g., [D, Chapter III.2.B], our current version of the theorem in [L-J] states that $[\mathfrak{a}]$ admits the factorization

$$[\mathfrak{a}] = \frac{1}{p!(-2\pi i)^p} d\varphi \circ R. \tag{1.3}$$

In particular, the right hand side is independent of the choice of bases of E_k.

If $\mathfrak{a}$ is generated by a regular sequence $f = (f_1, \dots, f_p)$ and (1.2) is the Koszul complex, then $d\varphi = p! df_p \wedge \cdots \wedge df_1$ and thus (1.3) reads

$$[\mathfrak{a}] = \frac{1}{(2\pi i)^p} R^f_{CH} \wedge df_p \wedge \cdots \wedge df_1, \tag{1.4}$$

cf. (1.1); this factorization appeared already in [CH], see also [DP]. A special case is the *Poincaré-Lelong formula*: if $f \in \mathcal{O}_0$, then $\bar\partial[1/f] \wedge df = 2\pi i [f = 0]$, where $[f = 0]$ is the current of integration along the zero set of f, counted with multiplicities.

The factorization (1.3) can be obtained from Lejeune-Jalabert's original result using residue theory. In a forthcoming paper [LW2] with Lärkäng we give a proof of (1.3), which does not rely on [L-J] and which extends to equi-dimensional ideals (i.e., all minimal primes are of the same dimension).

In [LW2] and [L-J], the factorization (1.3) is proved by comparing R and $d\varphi$ to a residue and differential form constructed from a certain Koszul complex; in [LW2] this is done using a recent comparison formula for residue currents due to Lärkäng, [L].

To explicitly describe the factors in (1.3) seems to be a delicate problem in general. In this note we compute the form $d\varphi$ when $\mathfrak{a}$ is a monomial ideal. More precisely, let A be the ring $\mathcal{O}_0$ of holomorphic germs at the origin in $\mathbf{C}^n$ with coordinates $z_1, \dots, z_n$, or let A be the polynomial ring $\mathbf{C}[z_1, \dots, z_n]$. We give an explicit description of the form

$$d\varphi = \sum_\sigma \frac{\partial \varphi_1}{\partial z_{\sigma(1)}} dz_{\sigma(1)} \circ \cdots \circ \frac{\partial \varphi_n}{\partial z_{\sigma(n)}} dz_{\sigma(n)} \tag{1.5}$$

when (1.2) is the *Scarf resolution*, introduced in [BPS], of an Artinian, i.e., zero-dimensional, *generic* monomial ideal M in A, see Section 3 for definitions. Here the sum is over all permutations σ of $\{1, \dots, n\}$. It turns out that each summand in (1.5) is a vector of monomials whose coefficients have a neat description in terms of the so-called staircase of M and sum up to the geometric multiplicity of M, see Theorem 1.1 below. This can be seen as a far-reaching generalization of the fact that the coefficent of dz^a equals a, which is the geometric multiplicity of the principal ideal (z^a), cf. Example 5.1 below. Thus in a certain sense the fundamental class of M is captured already in the form $d\varphi$. In the case of the Scarf resolution we recently, together with Lärkäng, [LW], gave an complete description of the current R. Combining Theorem 1.1 below with Theorem 1.1 in [LW] we obtain new proof of (1.3) in this case, cf. Corollary 1.2 below.

Let us describe our result in more detail. Let M be an Artinian monomial ideal in A. By the *staircase* $S = S_M$ of M we will mean the set

$$S = \overline{\{(x_1, \dots, x_n) \in \mathbf{R}^n_{>0} | z_1^{\lfloor x_1 \rfloor} \cdots z_n^{\lfloor x_n \rfloor} \notin M\}} \subset \mathbf{R}^n. \tag{1.6}$$

The name is motivated by the shape of S, cf. Figures 5.1, 5.2, and 6.1. We will refer to the finitely many maximal elements in S as *outer corners*.

The Scarf resolution $E_\bullet, \varphi_\bullet$ of M is encoded in the *Scarf complex*, Δ_M, which is a labeled simplicial complex of dimension $n-1$ with one vertex for each minimal monomial generator of M and one top-dimensional simplex for each outer corner of S, see Section 3. The rank of E_k equals the number of $(k-1)$-dimensional simplices in Δ_M. In particular, $E_\bullet, \varphi_\bullet$ ends at level n and the rank of E_n equals the number of outer cornes of S. Thus $d\varphi$ is a vector with one entry for each outer corner of S.

For our description of $d\varphi$ we need to introduce certain partitions of S. Given a permutation σ of $\{1, \dots, n\}$ let $\geq_\sigma$ be the lexicographical order induced by σ, i.e, $\alpha = (\alpha_1, \dots, \alpha_n) \geq_\sigma \beta = (\beta_1, \dots, \beta_n)$ if (and only if) for some $1 \leq k \leq n$, $\alpha_{\sigma(\ell)} = \beta_{\sigma(\ell)}$ for $\ell \leq k-1$ and $\alpha_{\sigma(k)} > \beta_{\sigma(k)}$, or $\alpha = \beta$. If $\alpha \neq \beta$ we write $\alpha >_\sigma \beta$. Let $\alpha^1 \geq_\sigma \dots \geq_\sigma \alpha^M$ be the total ordering of the outer corners induced by $\geq_\sigma$, and define inductively

$$\begin{aligned} S_{\sigma,\alpha^1} &= \{x \in S \mid x \leq \alpha_1\} \\ &\vdots \\ S_{\sigma,\alpha^k} &= \{x \in S \setminus (S_{\sigma,\alpha^1} \cup \dots \cup S_{\sigma,\alpha^{k-1}}) \mid x \leq \alpha_k\} \\ &\vdots \end{aligned}$$

For a fixed σ, $\{S_{\sigma,\alpha}\}_\alpha$ provides a partition of S, cf. Section 2.

Theorem 1.1. *Let M be an Artinian generic monomial ideal in A, and let*

$$0 \to E_n \xrightarrow{\varphi_n} \dots \xrightarrow{\varphi_2} E_1 \xrightarrow{\varphi_1} E_0 \tag{1.7}$$

be the Scarf resolution of A/M. Then

$$\frac{\partial \varphi_1}{\partial z_{\sigma(1)}} dz_{\sigma(1)} \circ \dots \circ \frac{\partial \varphi_n}{\partial z_{\sigma(n)}} dz_{\sigma(n)} = \sum_\alpha \operatorname{sgn}(\alpha) \operatorname{Vol}(S_{\sigma,\alpha}) z^{\alpha - \mathbf{1}} dz e^*_\alpha \tag{1.8}$$

where the sum is over the outer corners α of S, $\operatorname{sgn}(\alpha) = \pm 1$ comes from the orientation of the Scarf complex, $\{e^*_\alpha\}$ is a basis for E^*_n, $dz = dz_n \wedge \dots \wedge dz_1$, and $z^{\alpha-\mathbf{1}} = z_1^{\alpha_1 - 1} \cdots z_n^{\alpha_n - 1}$ if $\alpha = (\alpha_1, \dots, \alpha_n)$.

Theorem 1.1 in [LW] asserts that the residue current associated with the Scarf resolution equals

$$R = \sum_\alpha \operatorname{sgn}(\alpha) \bar\partial \left[\frac{1}{z_1^{\alpha_1}}\right] \circ \dots \circ \bar\partial \left[\frac{1}{z_n^{\alpha_n}}\right] e_\alpha, \tag{1.9}$$

where $\{e_\alpha\}$ is the basis of E_n dual to $\{e^*_\alpha\}$. Since $z^{a-1}\bar\partial[1/z^a] = [z=0]$, cf. (1.4), we conclude from (1.7) and (1.8) that

$$d_\sigma \varphi \circ R = (-2\pi i)^n \sum_\alpha \operatorname{Vol}(S_{\sigma,\alpha})[0] = (-2\pi i)^n \operatorname{Vol}(S)[0],$$

where $d_\sigma \varphi$ denotes the left hand side of (1.8). Note that $\operatorname{Vol}(S)$ equals the number of monomials that are not in M. Since these monomials form a basis for A/M, $\operatorname{Vol}(S) = \dim_{\mathbf{C}}(\mathcal{O}_0/M)$. Hence we get the following refinement of (1.3).

Corollary 1.2. *Let M and (1.7) be as in Theorem 1.1 and let R be the associated residue current. Then*

$$\frac{\partial \varphi_1}{\partial z_{\sigma(1)}} dz_{\sigma(1)} \wedge \cdots \wedge \frac{\partial \varphi_n}{\partial z_{\sigma(n)}} dz_{\sigma(n)} \circ R = (-2\pi i)^n [M].$$

Summing over all permutations σ we get back (1.3).

The core of the proof of Theorem 1.1 is an alternative description of the $S_{\sigma,\alpha}$ as certain cuboids, see Lemma 4.1. Given this description it is fairly straightforward to see that the volumes of $S_{\sigma,\alpha}$ are precisely the coefficents of the monomials in $d_\sigma \varphi$.

We suspect that Theorem 1.1 and Corollary 1.2 extend to a more general setting than the one above. Indeed, if M is an Artinian non-generic monomial ideal, we can still construct the partitions $\{S_{\sigma,\alpha}\}_\alpha$. The elements $S_{\sigma,\alpha}$ will, however, no longer be cuboids in general. Also, the computation of $d\varphi$ is more delicate in general. In Example 6.1 we compute $d\varphi$ for a non-generic monomial ideal for which the *hull resolution*, introduced in [BS], is minimal, and show that Theorem 1.1 holds in this case. On the other hand, in Example 6.2 we consider an example where the hull resolution is not a minimal resolution and where Theorem 1.1 fails to hold. However, for all examples we have considered Corollary 1.2 does hold, which raises the question whether it is always true that

$$\frac{\partial \varphi_1}{\partial z_{\sigma(1)}} dz_{\sigma(1)} \circ \cdots \circ \frac{\partial \varphi_n}{\partial z_{\sigma(n)}} dz_{\sigma(n)} \circ R = (-2\pi i)^p [\mathfrak{a}]. \tag{1.10}$$

We do not know whether the proofs of (1.3) in [LW2] and [L-J] could be modified to answer this question.

The paper is organized as follows. In Section 2 and 3 we provide some background on staircases of monomial ideals and the Scarf complex, respectively. In Section 3 we also present a version Theorem 3.1 of Theorem 1.1 where we specify the sign $\operatorname{sgn}(\alpha)$. The proof of Theorem 1.1 occupies Section 4 and in Section 5 we illustrate the proof by some examples. Finally, in Section 6 we consider resolutions of non-generic monomial ideals and look at some examples.

Acknowledgment. I want to thank Mats Andersson, Richard Lärkäng, and Jan Stevens for valuable discussions.

2. Staircases

We let $\geq$ denote the standard partial order on $\mathbf{R}^n$, i.e., $a = (a_1, \ldots, a_n) \geq b = (b_1, \ldots, b_n)$ if and only if $a_\ell \geq b_\ell$ for all ℓ. If $a_\ell > b_\ell$ for all ℓ we write $a \succ b$. Throughout we let A denote the ring $\mathbf{C}[z_1, \ldots, z_n]$ or the ring of holomorphic germs at $0 \in \mathbf{C}^n_{z_1,\ldots,z_n}$. For $a = (a_1, \ldots, a_n) \in \mathbf{N}^n$, where $\mathbf{N} = 0, 1, \ldots$, we will use the short hand notation z^a for the monomial $z_1^{a_1} \cdots z_n^{a_n}$ in A. For a general reference on (resolutions of) monomial ideals, see, e.g., [MS]

Unless otherwise stated M will be a monomial ideal in A, i.e., an ideal generated by monomials, and S will be the staircase of M as defined in (1.6). We let ∂S denote the boundary of S and and $S^c = \mathbf{R}^n_{>0} \setminus S$ the complement of S in $\mathbf{R}^n_{>0}$. As in the introduction we will refer to the maximal elements of S as *outer corners*. The minimal elements of the closure $\overline{S^c}$ of S^c in $\mathbf{R}^n$ we will call *inner corners*.

One can check that for any monomial ideal M there is a unique set of exponents $B \subset \mathbf{N}^n$ such that the monomials $\{z^a\}_{a \in B}$ generate M. We refer to these monomials as *minimal monomial generators* of M. Moreover

$$S = \mathbf{R}^n_{>0} \setminus \bigcup_{a\in B} (a + \mathbf{R}^n_{>0}). \tag{2.1}$$

In particular, the inner corners of S are precisely the elements in B.

Dually, M can be described as an intersection of so-called *irreducible* monomial ideals, i.e., ideals of the form $(z_1^{\alpha_1}, \dots, z_n^{\alpha_n}) =: \mathfrak{m}^\alpha$ (more generally an ideal is irreducible if it can not be written as a non-trivial intersection of two ideals). For every monomial ideal M there is a unique minimal set $C \subset \mathbf{N}^n$ such that

$$M = \bigcap_{\alpha\in C} \mathfrak{m}^\alpha. \tag{2.2}$$

The ideal M is Artinian if and only if each $\alpha \in C$ satisfies $\alpha \succ 0$ ($\alpha_\ell > 0$ for each ℓ). If $\alpha \succ 0$, then note that a monomial $z^b \notin \mathfrak{m}^\alpha$ if and only if $b \prec \alpha$. It follows that

$$S = \bigcup_{\alpha\in C} \{x \in \mathbf{R}^n_{>0} \mid x \le \alpha\}. \tag{2.3}$$

In particular, the outer corners of S are precisely the elements in C. If M is not Artinian, then S is not bounded in $\mathbf{R}^n$ and the representation (2.3) fails to hold. Note that (2.3) guarantees that for a fixed σ, $\{S_{\sigma,\alpha}\}_\alpha$, as defined in the introduction, is a partition of S.

Inspired by (2.1) we will call any set of this form a staircase. Let H be an affine subspace of $\mathbf{R}^n$ spanned by an affine sublattice L of $\mathbf{Z}^n$. For $a \in L$ let $U_a = U_a^L = \{x \in H \mid x \succ a\}$. Note that $U_0^{\mathbf{Z}^n}$ is just the first (open) orthant $\mathbf{R}^n_{>0}$ in $\mathbf{R}^n$. We say that a set $S \subset H$ is a *staircase* if it is of the form

$$S = U_{a^0} \setminus \bigcup_{j=1}^{s} U_{a^j}$$

for some $a^0, a^1, \dots, a^s \in L$. We say that a_0 is the *origin* of S and that $a^1, \dots, a^s$ are the *inner corners* of S. We call the maximal elements of S the *outer corners* of S. Since L is a lattice, the outer corners are in L.

Assume that L has rank k and that $\rho : L \to \mathbf{Z}^k$ is a lattice isomorphism that maps a^0 to $0 \in \mathbf{Z}^k$. Let $M(S)$ be the monomial ideal in $\mathbf{C}[z_1, \dots, z_k]$ (or the ring of holomorphic germs at $0 \in \mathbf{C}^k_{z_1,\dots,z_k}$) that is generated by y $z^{\rho(a^j)}$, where a^j are the inner corners of S. Then the staircase of $M(S)$ equals $\rho(S)$.

For $\alpha \in L$, let $V_\alpha = V_\alpha^L = \{x \in H \mid x \le \alpha\}$. If $M(S)$ is Artinian, then S admits a representation analogous to (2.3),

$$S = U_{a^0} \cap \bigcup_{\alpha} V_\alpha, \tag{2.4}$$

where the union is taken over all outer corners of S. In particular, S is a *lower set* in U_{a^0}: if $x \in S$, then $\{y \in U_{a^0} \mid y \le x\} \subset S$.

3. The Scarf complex

For $a, b \in \mathbf{R}^n$, we will denote by $a \vee b$ the *join* of a and b, i.e., the unique c such that $c \ge a, b$, and $c \le d$ for all $d \ge a, b$.

Let M be an Artinian monomial ideal in A, with minimal monomial generators $m_1 = z^{a^1}, \dots, m_r = z^{a^r}$. The *Scarf complex* $\Delta = \Delta_M$ of M was introduced by

Bayer-Peeva-Sturmfels, [BPS], based on previous work by H. Scarf. It is the collection of subsets $\mathcal{I} = \{i_1, \dots, i_k\} \subset \{1, \dots, r\}$ whose corresponding least common multiple $m_{\mathcal{I}} := \mathrm{lcm}(m_{i_1}, \dots, m_{i_k}) = z^{a^{i_1} \vee \cdots \vee a^{i_k}}$ is unique, that is,

$$\Delta = \{\mathcal{I} \subset \{1, \dots, r\} | m_{\mathcal{I}} = m_{\mathcal{I}'} \Rightarrow \mathcal{I} = \mathcal{I}'\}.$$

Clearly the vertices of Δ are the minimal monomial generators of M, i.e., the inner corners of S_M. One can prove that the Scarf complex is a simplicial complex of dimension is a most $n-1$. We let $\Delta(k)$ denote the simplices in Δ with k vertices, i.e., of dimension $k-1$. Moreover we will label the faces $\mathcal{I} \subset \Delta$ by the monomials $m_{\mathcal{I}}$.

We will sometimes be sloppy and identify the faces in Δ with their labels or exponents of the labels and write $m_{\mathcal{I}}$ or α for the face with label $m_{\mathcal{I}} = z^\alpha$. This means we will sometimes also write $\{z^{a^{i_1}}, \dots, z^{a^{i_k}}\}$ or $\{a^{i_1}, \dots, a^{i_k}\}$ for $\mathcal{I} = \{i_1, \dots, i_k\}$.

The ideal M is said to be *generic* in the sense of [BPS, MSY] if whenever two distinct minimal generators m_i and m_j have the same positive degree in some variable, then there is a third generator m_k that *strictly divides* $m_{\{i,j\}}$, which means that m_k divides $m_{\{i,j\}}/z_\ell$ for all variables z_ℓ dividing $m_{\{i,j\}}$. In particular, M is generic if no two generators have the same positive degree in any variable.

If M is generic, then Δ has precisely dimension $n-1$; it is a regular triangulation of the $(n-1)$-dimensional simplex. The (labels of) the top-dimensional faces of Δ are precisely the exponents α in a minimal irreducible decomposition (2.2) of M, i.e., the outer corners of S_M.

For $k = 0, \dots, n$, let E_k be the free A-module with basis $\{e_{\mathcal{I}}\}_{\mathcal{I} \in \Delta(k)}$ and let the differential $\varphi_k : E_k \to E_{k-1}$ be defined by

$$\varphi_k : e_{\mathcal{I}} \mapsto \sum_{j=1}^{k} (-1)^{j-1} \frac{m_{\mathcal{I}}}{m_{\mathcal{I}_j}} e_{\mathcal{I}_j}, \tag{3.1}$$

where $\mathcal{I}_j$ denotes $\{i_1, \dots, i_{j-1}, i_{j+1}, \dots, i_k\}$ if $\mathcal{I} = \{i_1, \dots, i_k\}$. Then the complex $E_\bullet, \varphi_\bullet$ is exact and thus gives a free resolution of the cokernel of φ_0, which with the identification $E_0 = A$ equals A/M, see [BPS, Theorem 3.2]. In fact, this so-called *Scarf resolution* is a minimal resolution of A/M, i.e., for each k, φ_k maps a basis of E_k to a minimal set of generators of $\mathrm{Im}\, \varphi_k$, see, e.g., [E, Corollary 1.5]. Originally, in [BPS], the situtation $A = \mathbf{C}[z_1, \dots, z_n]$ was considered. However, since $\mathcal{O}_0$ is flat over $\mathbf{C}[z_1, \dots, z_n]$, see, e.g., [T, Theorem 13.3.5], the complex $E_\bullet, \varphi_\bullet$ is exact for $A = \mathcal{O}_0$ if and only if it is exact for $A = \mathbf{C}[z_1, \dots, z_n]$.

Let $\mathcal{I} = \{i_1, \dots, i_n\}$ be a top-dimensional simplex in Δ with label α. Then there is a permutation η of $\{1, \dots, n\}$ such that for $1 \leq \ell \leq n$ there is a unique vertex $i_{\eta(\ell)}$ of $\mathcal{I}$ that satisfy $\alpha_\ell = a_\ell^{i_{\eta(\ell)}}$; we will refer to this vertex as the *x_ℓ-vertex* of $\mathcal{I}$. To see this, first of all, since $\alpha = \mathrm{lcm}(a^i)$, $a_\ell^i \leq \alpha_\ell$ and there must be at least one vertex i of $\mathcal{I}$ such that $a_\ell^i = \alpha_\ell$. Assume that i and j are vertices of $\mathcal{I}$ such that $a_\ell^i = a_\ell^j = \alpha_\ell$. Then, since M is generic, there is a generator z^b of M such that b strictly divides $a^i \vee a^j$. But then $a^i \vee a^j = a^i \vee a^j \vee b$ and so $\{i, j\}$ is not in Δ, which contradicts that i and j are both vertices of $\mathcal{I}$. We can now specify the sign in Theorem 1.1. Let $\mathrm{sgn}(\eta)$ denote the sign of the permutation η.

Theorem 3.1. *Let M and (1.2) be as in Theorem 1.1. Then, using the notation from Theorem 1.1,*

$$d_\sigma\varphi = \sum_\alpha \operatorname{sgn}(\eta)\operatorname{Vol}(S_{\sigma,\alpha}) z^{\alpha-\mathbf{1}} dz e^*_\alpha.$$

Also the sign in (1.9) equals $\operatorname{sgn}(\eta)$, see [LW, Theorem 5.1]. We should remark that we use a different sign convention in this paper than in [LW], which essentially corresponds to a different orientation of the $(n-1)$-simplex or to a different choice of bases for the modules E_k, cf. [AW, LW2]. In this paper we will use the convention that if (1.2) is the Koszul complex, then the coefficient of the only basis element of E_n is $\bar\partial[1/f_1]\wedge\cdots\wedge\bar\partial[1/f_n]$, whereas in [LW] it equals $\bar\partial[1/f_n]\wedge\cdots\wedge\bar\partial[1/f_1]$.

3.1. **The subcomplex $\Delta_{\sigma,a^1,\dots,a^k}$.** Given a permutation σ of $\{1,\dots,n\}$, and vertices $a^1,\dots,a^k$ of Δ, let $\Delta_{\sigma,a^1,\dots,a^k}$ be the (possibly empty) subcomplex of Δ, with top-dimensional simplices α that satisfy $\alpha_{\sigma(\ell)} = a^\ell_{\sigma(\ell)}$ for $\ell = 1,\dots,k$. In other words, the top-dimensional simplices in $\Delta_{\sigma,a^1,\dots,a^k}$ are the ones that have a^ℓ as $x_{\sigma(\ell)}$-vertex for $\ell = 1,\dots,k$.

Note that for each choice of permutation σ, $k\in\{1,\dots,n\}$, and $\alpha\in\Delta(n)$ there is a unique sequence $a^1,\dots,a^k$ such that $\alpha\in\Delta_{\sigma,a^1,\dots,a^k}$. Moreover, $\Delta_{\sigma,a^1,\dots,a^n}$ is the unique simplex $\alpha\in\Delta(n)$ that satisfies $a^\ell_{\sigma(\ell)} = \alpha_{\sigma(\ell)}$ for $\ell = 1,\dots,n$.

We will write $\Delta^*_{\sigma,a^1,\dots,a^k}$ for the subcomplex of $\Delta_{\sigma,a^1,\dots,a^k}$ consisting of all faces in $\Delta_{\sigma,a^1,\dots,a^k}$ that do not contain $a^1,\dots,a^{k-1}$, or a^k. Note that since $\Delta_{\sigma,a^1,\dots,a^k}$ is simplicial, $\{b^1,\dots,b^\ell\}$ is a face of $\Delta^*_{\sigma,a^1,\dots,a^k}$ if and only $\{a^1,\dots,a^k,b^1,\dots,b^\ell\}$ is a face of $\Delta_{\sigma,a^1,\dots,a^k}$.

4. Proof of Theorem 1.1

To prove the theorem we will first give an alternative description of the $S_{\sigma,\alpha}$ as certain cuboids. Througout this section we will assume that $E_\bullet,\varphi_\bullet$ is the Scarf resolution of a generic monomial ideal and we will use the notation of last section.

Lemma 4.1. *Assume that α is the label of the face $\mathcal{I} = \{i_1,\dots,i_n\}\in\Delta(n)$. Let η be the permutation of $\{1,\dots,n\}$ such that $i_{\eta(\ell)}$ is the x_ℓ-vertex of α and set $\tau = \eta\circ\sigma$.*

Then $S_{\sigma,\alpha}$ is a cuboid with side lengths

$$a^{i_{\tau(1)}}_{\sigma(1)}, \left(a^{i_{\tau(1)}}\vee a^{i_{\tau(2)}} - a^{i_{\tau(1)}}\right)_{\sigma(2)}, \dots, \left(a^{i_{\tau(1)}}\vee\cdots\vee a^{i_{\tau(n)}} - a^{i_{\tau(1)}}\vee\cdots\vee a^{i_{\tau(n-1)}}\right)_{\sigma(n)}.$$

In particular,

$$\operatorname{Vol}(S_{\sigma,\alpha}) = a^{i_{\tau(1)}}_{\sigma(1)}\times\left(a^{i_{\tau(1)}}\vee a^{i_{\tau(2)}} - a^{i_{\tau(1)}}\right)_{\sigma(2)}\times\cdots\times\left(a^{i_{\tau(1)}}\vee\cdots\vee a^{i_{\tau(n)}} - a^{i_{\tau(1)}}\vee\cdots\vee a^{i_{\tau(n-1)}}\right)_{\sigma(n)}.$$

4.1. **Proof of Lemma 4.1.** To prove the lemma, let us first assume that σ is the identity permutation and write $S_\alpha = S_{\sigma,\alpha}$, $\Delta_{a^1,\dots,a^k} = \Delta_{\sigma,a^1,\dots,a^k}$, and $\Delta^*_{a^1,\dots,a^k} = \Delta^*_{\sigma,a^1,\dots,a^k}$.

We will decompose S in a seemingly different way. First we will construct certain lower-dimensional staircases with corners in Δ. Given an inner corner $a = (a_1,\dots,a_n)$ of S let

$$R_a = \{x\in S \mid x_1 = a_1, x_\ell > a_\ell, \ell = 2,\dots,n\}.$$

Note that $R_a\subset\partial S$.

Claim 4.2. *Let S be the staircase in $\mathbf{R}^n_{x_1,\dots,x_n}$ of an Artinian generic monomial ideal, and let a be an inner corner of S. Then R_a is a staircase in the hyperplane $\{x_1 = a_1\}$ with origin a. The outer corners of R_a are the top-dimensional faces of Δ_a. The inner corners are the lattice points $a \vee b$, where b is a vertex in Δ_a^*.*

In particular R_a is empty precisely if Δ_a^* is empty.

Proof. Assume that $\beta = (\beta_1, \dots, \beta_n)$ is maximal in R_a. Since $R_a \subset \{x_1 = a_1\}$, $\beta_1 = a_1$. Moreover, since S is a lower set, there is a $\gamma \geq \beta$ that is maximal in S. Assume that $\gamma_1 > a_1$. By the definition of R_a, $\gamma_\ell \geq \beta_\ell > a_\ell$ for $\ell = 2, \dots, n$ and thus $\gamma \succ a$, which contradicts that $\gamma \in S$. Hence $\gamma_1 = a_1$ and since $\gamma \geq \beta \geq a$, $\gamma \in R_a$ and thus $\gamma = \beta$, so, in fact, β is maximal in S. Since $\beta_1 = a_1$, we conclude that $\beta \in \Delta_a(n)$.

On the other hand, if $\beta \in \Delta_a(n)$, then β is maximal in S and contained in R_a, and thus is maximal in R_a. Thus we conclude that the maximal elements in R_a are the top-dimensional faces of Δ_a.

Since S is a lower set,

$$R_a = \{x \mid x_1 = a_1, x_\ell > a_\ell, \ell = 2, \dots, n\} \cap \bigcup_{\beta \in \Delta_a(n)} \{x \mid x \leq \beta\},$$

which in light of (2.4) means that R_a is a staircase in $\{x_1 = a_1\}$ with origin a and outer corners $\Delta_a(n)$.

Next, assume that $\beta = (a_1, \beta_2, \dots, \beta_n)$ is minimal in $\overline{R_a^c} := \overline{S^c} \cap \{x_1 = a_1\}$. Then there is a minimal $b = (b_1, \dots, b_n) \leq \beta$ in $\overline{S^c}$. Assume that $b_j < \beta_j$ for some $j = 2, \dots, n$. Then $(a \vee b) \in \overline{R_a^c}$ and $a \vee b < \beta$, which contradicts that β is minimal in $\overline{R_a^c}$. Thus we can conclude that $b_\ell = \beta_\ell$ for $\ell = 2, \dots, n$; in particular, $\beta = a \vee b$. Since β is minimal in $\overline{R_a^c}$, β and thus b divide at least one $\alpha \in \Delta_a(n)$. It follows that $b \in \Delta_a^*(1)$.

Conversely, pick $b \in \Delta_a^*(1)$. Then $\beta := a \vee b \in \overline{S^c} \cap \{x_1 = a_1\} = \overline{R_a^c}$. Assume that β is not minimal in $\overline{R_a^c}$. Then there is a $\gamma = (a_1, \gamma_2, \dots, \gamma_n) \neq \beta$ such that $\gamma \leq \beta$, and as above a minimal $c \leq \gamma$ in $\overline{S^c}$ of the form $c = (c_1, \gamma_2, \dots, \gamma_n)$. Now $\operatorname{lcm}(z^a, z^b, z^c) = \operatorname{lcm}(z^a, z^b)$, which contradicts that $\{a, b\} \in \Delta_a \subset \Delta$. Hence β is minimal in $\overline{R_a^c}$. We conclude that the inner corners of R_a are exactly the lattice points $a \vee b$, where $b \in \Delta_a^*(1)$.

□

Let $\pi : \mathbf{R}^n_{x_1,\dots,x_n} \to \mathbf{R}^{n-1}_{x_2,\dots,x_n}$ be the projection $\pi : (x_1, \dots, x_n) \mapsto (x_2, \dots, x_n)$ and let $\rho : \mathbf{R}^n_{x_1,\dots,x_n} \to \mathbf{R}^{n-1}_{x_2,\dots,x_n}$ be the affine mapping defined by $\rho(x) = \pi(x - a)$. Then the restriction to $\{x_1 = a_1\} \cap \mathbf{Z}^n$ is a lattice isomorphism that maps a to $0 \in \mathbf{Z}^{n-1}$. Let M_a be the monomial ideal in $\mathbf{C}[z_2, \dots, z_n]$ (or the ring of holomorphic germs at $0 \in \mathbf{C}^{n-1}_{z_2,\dots,z_n}$) defined by R_a and ρ as in Section 2.

Note that Claim 4.2 implies in particular that if for $b \in \Delta_a(1)$, $(b_2, \dots, b_n) \geq (a_2, \dots, a_n)$ and thus $\rho(a \vee b) = \rho(b)$. Thus, in fact, M_a is generated by $\rho(b)$, where b is a vertex of Δ_a^*.

Claim 4.3. *M_a is a generic Artinian monomial ideal.*

Proof. First, M_a is Artinian since R_a is bounded.

To show that M_a is generic, assume that β and γ are minimial generators of M_a such that $\beta_\ell = \gamma_\ell$ for some $2 \leq \ell \leq n$. Then $\beta = \rho(b)$ and $\gamma = \rho(c)$ for some minimal generators b and c of M. Note that $\beta_\ell = \gamma_\ell$ is equivalent to $b_\ell = c_\ell$. Since M is generic there exists a minimal generator z^d of M that strictly divides $\operatorname{lcm}(z^b, z^c)$, i.e., $d_\ell < (b \vee c)_\ell$ for all ℓ such that $(b \vee c)_\ell > 0$. Set $\delta = \rho(d)$. It follows that $\delta_\ell < (\beta \vee \gamma)_\ell$ for all ℓ such that $(\beta \vee \gamma)_\ell > 0$, so that z^δ strictly divides $\operatorname{lcm}(z^\beta, z^\gamma)$.

Assume that $z^\delta \notin M_a$. Then there is an outer corner of R_a, i.e., $\alpha \in \Delta_a(n)$, with vertices a and, say, $a^2, \dots, a^n$, such that $\delta \leq \rho(\alpha)$. It follows that $\operatorname{lcm}(z^\delta, z^a, z^{a^2}, \dots, z^{a^n}) = \operatorname{lcm}(z^a, z^{a^2}, \dots, z^{a^n})$, which contradicts that $\alpha \in \Delta$. Therefore $z^\delta \in M_a$ and we conclude that M_a is generic. □

Claim 4.4. *The Scarf complex Δ_{M_a} of M_a equals Δ_a^* but with a different labeling. More precisely, Δ_{M_a} consists exactly of the faces of the form $\{\rho(b^1), \dots, \rho(b^j)\}$, where $\{b^1, \dots, b^j\}$ is a face of Δ_a^*.*

Proof. We have seen above that the minimal generators of M_a are precisely the monomials of the form $z^{\rho(b^\ell)}$, where b^ℓ is a vertex of Δ_a^*. Thus it suffices to prove that the least common multiple of $\rho(z^{b^1}), \dots, \rho(z^{b^j})$ is unique if and only if the least common multiple of $z^{b^1}, \dots, z^{b^j}$ is unique, or equivalently that $\rho(b^1) \vee \cdots \vee \rho(b^j)$ is unique if and only if $b^1 \vee \cdots \vee b^j$ is unique, when b^j are among the vertices of Δ_a^*. Now, since ρ is a lattice isomorphism, $\rho(b^1) \vee \cdots \vee \rho(b^j) = \rho(a \vee b^1) \vee \cdots \vee \rho(a \vee b^j)$ is unique if and only if $(a \vee b^1) \vee \cdots \vee (a \vee b^j) = a \vee b^1 \vee \cdots \vee b^j$ is unique. However, since $b^\ell \in \Delta_a^*(0)$, $a \vee b^1 \vee \cdots \vee b^j$ is unique if and only if $b^1 \vee \cdots \vee b^j$ is unique.

□

Claim 4.5. *Assume that $a \neq b$ are inner corners of S. Then $\pi(R_a) \cap \pi(R_b) = \emptyset$.*

Proof. We first assume that $a_1 \neq b_1$ and that $\pi(R_a) \cap \pi(R_b) \neq \emptyset$; pick $\beta \in \pi(R_a) \cap \pi(R_b)$. Without loss of generality we may assume that $a_1 > b_1$. Since $\beta \in \pi(R_b)$, $\beta_j > b_j$ for $j = 2, \dots, n$ and thus $\gamma := (a_1, \beta_2, \dots, \beta_n)$ is contained in the interior of S^c. On the other hand, $\beta \in \pi(R_a)$ implies that $\gamma \in R_a \subset \partial S$. Thus $\pi(R_a) \cap \pi(R_b) = \emptyset$ if $a_1 \neq b_1$.

Next, assume that $a_1 = b_1$ and that $\pi(R_a) \cap \pi(R_b) \neq \emptyset$. Since R_a and R_b are both contained in the hyperplane $\{x_1 = a_1\}$, $\pi(R_a) \cap \pi(R_b) \neq \emptyset$ is equivalent to that $R_a \cap R_b \neq \emptyset$. Assume that $\beta \in R_a \cap R_b$. Since M is generic there is a minimal generator z^c that strictly divides $\operatorname{lcm}(z^a, z^b)$. It follows that $\beta_\ell > c_\ell$ for $\ell = 1, \dots, n$, and thus β is contained in the interior of S^c, which contradicts that $\beta \in R_a \cap R_b$. Thus we have proved that $\pi(R_a) \cap \pi(R_b) = \emptyset$ when $a \neq b$. □

Claim 4.6. *For each $x \in S$, there is an inner corner a of S such that $\pi(x) \in \pi(R_a)$.*

Proof. Assume that $x \in S$. Then there is a minimal generator z^a, such that $\pi(x)_\ell > \pi(a)_\ell$ for $\ell = 2, \dots, n$. Indeed, since M is Artinian, there is a minimal generator of the form $z_1^{a_1}$, whose exponent is mapped to the origin in $\mathbf{R}^{n-1}$.

On the other hand, since M is Artinian, if $\pi(b)$ is maximal among the images $\pi(a)$ of inner corners a, then in fact the x_1-coordinate of b is zero, which implies that $\pi(x)_\ell$ cannot be greater than $\pi(b)_\ell$ for all $\ell = 2, \dots, n$.

Now the claim follows from the following fact, which is a consequence of Claim 4.2: each $\xi \in \mathbf{R}^{n-1}$, such that $\xi_\ell > \pi(a)_\ell$ for $\ell = 2, \dots, n$ is in $\pi(R_a)$ unless $\xi_\ell > \pi(b)_\ell$ for $\ell = 2, \dots, n$ for some inner corner b such that $\pi(b) \geq \pi(a)$.

□

Next, we will use the staircases R_a to construct a partition of S. For each inner corner of S, let

$$P_a = \{x \in S \mid \pi(x) \in \pi(R_a)\}.$$

In other words, P_a consists of everything in S "below" the staircase R_a. By a slight abuse of notation, $P_a =]0, a_1] \times R_a$. From Claims 4.5 and 4.6 it follows that the (non-empty) P_a give a partition of S. Note that P_a is a staircase itself with origin $\pi(a)$, outer corners $\alpha \in \Delta_a(n)$, and inner corners a and $\pi(b)$, where b is a vertex of Δ_a^* and where we think of $\pi(\mathbf{R}^n)$ as embedded as the hyperplane $\{x_1 = 0\}$ in $\mathbf{R}^n$.

Claim 4.7. *If* $\alpha \in \Delta(n)$ *is contained in* Δ_a*, then* $S_\alpha \subset P_a$*.*

Proof. First, note that

$$S_\alpha = \{x \leq \alpha\} \cap \Big(S \setminus \bigcup_{\beta >_\sigma \alpha} S_\beta\Big) \subset \{x \leq \alpha\} \cap \Big(S \setminus \bigcup_{\beta_1 > \alpha_1} S_\beta\Big). \tag{4.1}$$

Next, let β be an outer corner of S such that $\beta_1 > \alpha_1$. Assume that $x \in P_a$ satisfies $x \leq \beta$. Then $\beta_\ell \geq x_\ell > a_\ell$ for $\ell = 2, \ldots, n$. Hence $\beta_\ell > a_\ell$ for $\ell = 1, \ldots, n$, which contradicts that a is an inner corner of S. Therefore $S_\beta \cap P_a = \emptyset$, and hence $P_a \subset \big(S \setminus \bigcup_{\beta_1 \geq \alpha_1} S_\beta\big)$. If $\alpha \in \Delta_a$, then from Claim 4.2 we know that α is contained in P_a and since P_a is a lower set we conclude that the rightmost expression in (4.1) is contained in P_a; in particular, $S_\alpha \subset P_a$.

□

Next, we will inductively define staircases and partitions associated with faces of Δ of higher dimension. Given vertices $a^1, \ldots, a^{k-1}$ of Δ, (such that $\Delta_{a^1,\ldots,a^{k-1}}$ is nonempty) and an inner corner a^k of $\Delta^*_{a^1,\ldots,a^{k-1}}$, assuming that $R_{a^1,\ldots,a^{k-1}}$ is defined, we let

$$R_{a^1,\ldots,a^k} := \{x \in R_{a^1,\ldots,a^{k-1}} \mid x_k = (a^1 \vee \cdots \vee a^k)_k, x_j > (a^1 \vee \cdots \vee a^k)_j, j = k+1, \ldots, n\}.$$

Let $\pi_k : \mathbf{R}^n_{x_1,\ldots,x_n} \to \mathbf{R}^{n-k}_{x_{k+1},\ldots,x_n}$ be the projection $\pi_k : (x_1, \ldots, x_n) \mapsto (x_{k+1}, \ldots, x_n)$, and let $\rho_k : \mathbf{R}^n_{x_1,\ldots,x_n} \to \mathbf{R}^{n-k}_{x_{k+1},\ldots,x_n}$ be the affine mapping defined by $\rho_k : x \mapsto \pi_k(x - a^1 \vee \cdots \vee a^k)$. Then note that the restriction to $H_k \cap \mathbf{Z}^n$, where H_k is the codimension k-plane

$$H_k = \{x_1 = (a^1 \vee \cdots \vee a^k)_1, \ldots, x_k = (a^1 \vee \cdots \vee a^k)_k\}$$

is a lattice isomorphism that maps $a^1 \vee \cdots \vee a^k$ to $0 \in \mathbf{Z}^{n-k}$. Note, in light of Section 3.1, that in fact $H_k = \{x_1 = a^1_1, \ldots, x_k = a^k_k\}$. Moreover, let $\iota_k : \mathbf{R}^{n-k}_{x_{k+1},\ldots,x_n} \to \mathbf{R}^n_{x_1,\ldots,x_n}$ be the embedding $\iota_k : (x_{k+1}, \ldots, x_n) \mapsto (0, \ldots, 0, x_{k+1}, \ldots, x_n)$. We will somewhat abusively denote by ρ_k^{-1} the mapping $\rho_k^{-1} : \mathbf{R}^{n-k}_{x_{k+1},\ldots,x_n} \to \mathbf{R}^n_{x_1,\ldots,x_n}$ that maps x to $\iota_k(x) + a^1 \vee \cdots \vee a^k$. The notation is motivated by the fact that $\rho_k^{-1} \circ \rho_k$ is the identity on H_k and $\rho_k \circ \rho_k^{-1}$ is the identity on $\mathbf{R}^{n-k}$.

Claim 4.8. *Assume that* $a^1, \ldots, a^k$ *are inner corners of* S*, such that* a^j *is in* $\Delta_{a^1,\ldots,a^{j-1}}$ *for* $j = 2, \ldots, k$*. Then* $R_{a^1,\ldots,a^k}$ *is a staircase in* H_k*. The origin of* $R_{a^1,\ldots,a^k}$ *is* $a^1 \vee \cdots \vee a^k$*, the outer corners are the top-dimensional faces of* $\Delta_{a^1,\ldots,a^k}$ *and the inner corners are the lattice points of the form* $a^1 \vee \ldots \vee a^k \vee b$*, where* b *is a vertex of* $\Delta^*_{a^1,\ldots,a^k}$*.*

The monomial ideal $M_{a^1,\dots,a^k}$ *defined by* $R_{a^1,\dots,a^k}$ *and* ρ_k *as in Section 2 is an Artinian generic monomial ideal. The Scarf complex of* $M_{a^1,\dots,a^k}$ *is equal to* $\Delta^*_{a^1,\dots,a^k}$*, with the face with label* α *relabeled by* $\rho_k(\alpha) = \rho_k(a^1 \vee \cdots \vee a^k \vee \alpha)$*.*

Proof. First, it is clear from the definition that $R_{a^1,\dots,a^k}$ is contained in H_k.

Assume that the claim is true for $k-1$.

To show that $R_{a^1,\dots,a^k}$ is a staircase, apply Claim 4.2 to the staircase $\widetilde{S} := \rho_{k-1}(R_{a^1,\dots,a^{k-1}})$ of $M_{a^1,\dots,a^{k-1}}$ in $\mathbf{R}^{n-k+1}_{x_k,\dots,x_n}$ and the inner corner $\tilde{a} := \rho_{k-1}(a^1 \vee \cdots \vee a^k)$ of $\widetilde{S}$. We get that

$$R_{\tilde{a}} = \{x \in \widetilde{S} \mid x_k = \tilde{a}_k, x_\ell > \tilde{a}_\ell, \ell = k+1,\dots,n\}$$

is a staircase in the hyperplane $\{x_k = \tilde{a}_k\}$ with origin $\tilde{a}$. The outer corners of $R_{\tilde{a}}$ are the top-dimensional faces of the subcomplex $\Delta_{\tilde{a}}$ of $\Delta_{M_{a^1,\dots,a^{k-1}}}$, which consists of the top-dimensional faces of $\Delta_{M_{a^1,\dots,a^{k-1}}}$ whose x_k-vertex is $\tilde{a}$. The inner corners of $R_{\tilde{a}}$ are the lattice points $\tilde{a} \vee \tilde{b}$ where $\tilde{b}$ is a vertex of $\Delta_{\tilde{a}}$ not equal to $\tilde{a}$.

Note that $\rho^{-1}_{k-1}(R_{\tilde{a}}) = R_{a^1,\dots,a^k}$. Moreover, note that $\tilde{\alpha}$ is a top-dimensional simpex in $\Delta_{\tilde{a}}$ if and only if $\rho^{-1}_{k-1}(\tilde{\alpha})$ is a top-dimensional simplex of $\Delta_{a^1,\dots,a^k}$, and $\tilde{b} := \rho_{k-1}(a^1 \vee \cdots \vee a^k \vee b)$ is a vertex of $\Delta_{\tilde{a}}$ if and only if b is a vertex of $\Delta_{a^1,\dots,a^k}$.

It follows that $R_{a^1,\dots,a^k}$ is a staircase in $\mathbf{R}^n$ with origin $\rho^{-1}_{k-1}(\tilde{a}) = a^1 \vee \cdots \vee a^k$, outer corners $\alpha \in \Delta_{a^1,\dots,a^k}(n)$ and inner corners $\rho^{-1}_{k-1}(\tilde{a} \vee \tilde{b}) = a^1 \vee \cdots \vee a^k \vee b$, where b is a vertex of $\Delta^*_{a^1,\dots,a^k}$.

Let $\rho' : \mathbf{R}^{n-k+1}_{x_k,\dots,x_n} \to \mathbf{R}^{n-k}_{x_{k+1},\dots,x_n}$ be the affine map $\rho' : (x_k,\dots,x_n) \mapsto (x_{k+1} - \tilde{a}_{k+1},\dots,x_n - \tilde{a}_n)$. Then the restriction to $\{x_k = \tilde{a}_k\} \cap \mathbf{Z}^{n-k+1}$ is a lattice isomorphism that maps $\tilde{a}$ to $0 \in \mathbf{Z}^{n-k}$. Note that $\rho_k = \rho' \circ \rho_{k-1}$. It follows that $M_{a^1,\dots,a^k}$ is the ideal defined by $R_{\tilde{a}}$ and ρ' as in Section 2 and thus by Claim 4.3 it is an Artinian generic monomial ideal. Moreover the generators of $M_{a^1,\dots,a^k}$ are precisely $\rho'(\tilde{a} \vee \tilde{b}) = \rho_k(a^1 \vee \cdots \vee a^k \vee b)$, where $b \in \Delta_{a^1,\dots,a^k}(1)$. Recall from the text after Claim 4.2 that $\rho'(\tilde{a} \vee \tilde{b}) = \rho'(\tilde{b})$. Thus

$$\rho_k(a^1 \vee \cdots \vee a^k \vee b) = \rho'(\tilde{b}) = \rho'(\rho_{k-1}(a^1 \vee \cdots \vee a^{k-1} \vee b)) = \rho'(\rho_{k-1}(b)) = \rho_k(b)$$

where we have used the induction hypothesis that the claim holds for $k-1$ for the third equality. By arguing as in the proof of Claim 4.4 one sees that $\rho_k(b^1) \vee \cdots \vee \rho_k(b^j)$ is unique if and only if $b^1 \vee \cdots \vee b^j$ is unique. Thus the Scarf complex of $M_{a^1,\dots,a^k}$ is of the desired form.

□

To construct the partitions associated with the staircases $R_{a^1,\dots,a^k}$, we define inductively

$$P_{a^1,\dots,a^k} = \{x \in P_{a^1,\dots,a^{k-1}} \mid \pi_k(x) \in \pi_k(R_{a^1,\dots,a^k})\}.$$

Then $P_{a^1,\dots,a^k}$ is a k-dimensional cuboid times the $(n-k)$-dimensional staircase $R_{a^1,\dots,a^k}$. The ℓth side length is given as the "height" of $R_{a^1,\dots,a^\ell}$ in $R_{a^1,\dots,a^{\ell-1}}$. By a slight abuse of notation

$$P_{a^1,\dots,a^k} =]0, a^1_1] \times]a^1_2, (a^1 \vee a^2)_2] \times \cdots \times](a^1 \vee \cdots \vee a^{k-1})_k, (a^1 \vee \cdots \vee a^k)_k] \times R_{a^1,\dots,a^k}.$$

In particular,

$$P_{a^1,\dots,a^n} =]0, a^1_1] \times]a^1_2, (a^1 \vee a^2)_2] \times \cdots \times](a^1 \vee \cdots \vee a^{n-1})_n, (a^1 \vee \cdots \vee a^n)_n]. \tag{4.2}$$

Note that $P_{a^1,\dots,a^k}$ is in fact a staircase with origin $\iota_k(\pi_k(a^1\vee\cdots\vee a^k))$, outer corners $\alpha\in\Delta_{a^1,\dots,a^k}(n)$, and inner corners $a^1\vee\cdots\vee a^k$ and $\iota_k(\pi_k(b))$, where $b\in\Delta^*_{a^1,\dots,a^k}(1)$.

By inductively applying Claims 4.5 and 4.6 we get that for each k the set of (nonvanishing) $P_{a^1,\dots,a^k}$ gives a partition of S.

Claim 4.9. *If $\alpha\in\Delta(n)$ is contained in $\Delta_{a^1,\dots,a^k}$, then $S_\alpha\subset P_{a^1,\dots,a^k}$.*

Proof. First note that

$$S_\alpha=\{x\le\alpha\}\cap\Big(S\setminus\bigcup_{\beta>_\sigma\alpha}S_\beta\Big)\subset\{x\le\alpha\}\cap\Big(S\setminus\bigcup_{j=1}^{k}\ \bigcup_{\beta_1=\alpha_1,\dots,\beta_{j-1}=\alpha_{j-1},\beta_j>\alpha_j}S_\beta\Big). \tag{4.3}$$

Next, assume that $\alpha\in\Delta_{a^1,\dots,a^k}(n)$ and pick $j\le k$. Let $a:=a^1\vee\cdots\vee a^j$, and let β be an outer corner of S such that $\beta_1=a_1,\dots,\beta_{j-1}=a_{j-1}$, and $\beta_j>a_j$. Assume that $x\in P_{a^1,\dots,a^j}$ satisfies $x\le\beta$. Then $\beta_\ell\ge x_\ell>a_\ell$ for $\ell=j+1,\dots,n$, so that $\beta_\ell>a_\ell$ for $\ell=j,\dots,n$. But this cannot happen since a is an inner corner of $R_{a^1,\dots,a^{j-1}}$. Thus

$$\emptyset=S_\beta\cap P_{a^1,\dots,a^j}\supset S_\beta\cap P_{a^1,\dots,a^k}$$

and it follows that

$$P_{a^1,\dots,a^k}\subset S\setminus\Big(\bigcup_{j=1}^{k}\ \bigcup_{\beta_1=\alpha_1,\dots,\beta_{j-1}=\alpha_{j-1},\beta_j>\alpha_j,j\le k}S_\beta\Big).$$

Since $\alpha\in P_{a^1,\dots,a^k}$ and $P_{a^1,\dots,a^k}$ is a staircase and thus a lower set we conclude that the rightmost expression in (4.3) is contained in $P_{a^1,\dots,a^k}$ and thus $S_\alpha\subset P_{a^1,\dots,a^k}$. ☐

Recall from Section 3.1 that $\Delta_{a^1,\dots,a^n}$ is just the simplex $\alpha\in\Delta(n)$ with vertices $a^1,\dots,a^n$. On the other hand, each outer corner α gives rise to a nonempty $P_\alpha:=P_{a^1,\dots,a^n}$ by choosing a^ℓ as the x_ℓ-vertex of α, cf. Section 3. By Claim 4.9, $S_\alpha\subset P_\alpha$ and since both P_α and S_α give partitions of S, we conclude that $S_\alpha=P_\alpha$.

Now given $\mathcal{I}=\{i_1,\dots,i_n\}\in\Delta(n)$ we choose a^ℓ as the x_ℓ-vertex $a^{i_{\eta(\ell)}}$. Then Lemma 4.1 follows in light of (4.2).

For a general choice of σ the above proof works verbatim, with coordinates x_ℓ and variables z_ℓ replaced by $x_{\sigma(\ell)}$ and $z_{\sigma(\ell)}$, respectively, and η replaced by τ.

4.2. **Computing $d\varphi$.** Let us now compute the $e^*_{\mathcal{I}}$-entry of $d_\sigma\varphi$ for a given $\mathcal{I}=\{i_1,\dots,i_n\}\in\Delta(n)$. Recall from (3.1) that

$$\varphi_k=\sum_{\mathcal{I}'=\{i'_1,\dots,i'_k\}\subset\mathcal{I}}(-1)^{j-1}z^{a^{i'_1}\vee\cdots\vee a^{i'_k}-a^{i'_1}\vee\cdots\widehat{a^{i'_j}}\cdots\vee a^{i'_k}}e^*_{\mathcal{I}'}\otimes e_{\mathcal{I}'_j}+\varphi'_k \tag{4.4}$$

where φ_k' are the remaining terms that will not contribute to the $e_{\mathcal{I}}^*$-entry. It follows that the coefficient of $e_{\mathcal{I}}^*$ in $d_\sigma\varphi$ equals

$$\begin{aligned}
&\operatorname{sgn}\big((n,\dots,1)\big)\sum_\tau \operatorname{sgn}(tau)\frac{\partial}{\partial z_{\sigma(1)}} z^{a^{i_{\tau(1)}}} dz_{\sigma(1)} \wedge \frac{\partial}{\partial z_{\sigma(2)}} z^{a^{i_{\tau(1)}}\vee a^{i_{\tau(2)}}-a^{i_{\tau(1)}}} dz_{\sigma(2)} \\
&\qquad \wedge\cdots\wedge \frac{\partial}{\partial z_{\sigma(n-1)}} z^{a^{i_{\tau(1)}}\vee\dots\vee a^{i_{\tau(n-1)}}-a^{i_{\tau(1)}}\vee\dots\vee a^{i_{\tau(n-2)}}} dz_{\sigma(n-1)}\wedge \\
&\qquad\qquad \frac{\partial}{\partial z_{\sigma(n)}} z^{a^{i_{\tau(1)}}\vee\dots\vee a^{i_{\tau(n)}}-a^{i_{\tau(1)}}\vee\dots\vee a^{i_{\tau(n-1)}}} dz_{\sigma(n)} =: \sum_\tau F_\tau,
\end{aligned} \tag{4.5}$$

where the sum is over all permutations τ of $\{1,\dots,n\}$.

Note that $z^{a^{i_{\tau(1)}}\vee\dots\vee a^{i_{\tau(k)}}-a^{i_{\tau(1)}}\vee\dots\vee a^{i_{\tau(k-1)}}}$ is a monomial in the variables $z_{\eta^{-1}(\tau(k))},\dots,z_{\eta^{-1}(\tau(n))}$. Therefore the last factor in F_τ vanishes unless $\tau(n)=\eta(\sigma(n))$. Given, $\tau(n)=\eta(\sigma(n))$, the next to last factor vanishes unless $\tau(n-1)=\eta(\sigma(n-1))$ etc. To conclude, F_τ, where $\tau=\eta\circ\sigma$, is the only nonvanishing term in (4.5).

Now with $\tau=\eta\circ\sigma$,

$$\begin{aligned}
F_\tau &= \operatorname{sgn}(\tau)\times a^{i_{\tau(1)}}_{\sigma(1)} \times \Big(a^{i_{\tau(1)}}\vee a^{i_{\tau(2)}} - a^{i_{\tau(1)}}\Big)_{\sigma(2)} \times\cdots\times \\
&\Big(a^{i_{\tau(1)}}\vee\cdots\vee a^{i_{\tau(n)}} - a^{i_{\tau(1)}}\vee\cdots\vee a^{i_{\tau(n-1)}}\Big)_{\sigma(n)} z^{a^{i_{\tau(1)}}\vee\dots\vee a^{i_{\tau(n)}}} \frac{dz_{\sigma(n)}}{z_{\sigma(n)}}\wedge\cdots\wedge\frac{dz_{\sigma(1)}}{z_{\sigma(1)}} = \\
&\qquad \operatorname{sgn}(\eta)\operatorname{Vol}(S_{\sigma,\alpha})z^{\alpha-\mathbf{1}}dz_n\wedge\cdots\wedge dz_1,
\end{aligned} \tag{4.6}$$

where the last equality follows from Lemma 4.1. This concludes the proof of Theorem 1.1.

5. Examples

Let us illustrate Theorem 1.1 by some examples.

Example 5.1. Assume that $n=1$. Then each monomial ideal M is a principal ideal generated by a monomial z^a. The staircase of M is just the line segment $]0,a]\subset\mathbf{R}_{>0}$ with one outer corner $\alpha=a$ so that $S_\alpha=S$. Moreover, the Scarf complex is just a point with label z^a, and thus the Scarf resolution is just $0\to A\xrightarrow{z^a}A$. Thus in this case Theorem 1.1 just reads

$$d\varphi = d(z^a) = \operatorname{Vol}(]0,a])z^{a-1}dz = az^{a-1}dz.$$

□

Example 5.2. Assume that $n=2$. Then each Artinian monomial ideal is of the form $M=(z_1^{a_1}z_2^{b_1},\dots,z_1^{a_r}z_2^{b_r})$ for some integers $a_1>\ldots>a_r=0$ and $0=b_1<\ldots<b_r$. Since no two minimal monomial generators have the same positive degree in any variable, M is trivially generic. In this case the staircase of M looks like a actual staircase with inner corners (a_j,b_j) and outer corners $\alpha^j:=(a_j,b_{j+1})$, see Figure 5.1. Note that $\sigma=(1,2)$ corresponds to the ordering $\alpha^1\geq_\sigma\ldots\geq_\sigma\alpha^{r-1}$ of the outer corners and

$$S_{(1,2),\alpha^j}=\{x\in\mathbf{R}^2_{>0}\mid 0\leq x_1<a_j, b_j\leq x_2<b_{j+1}\},$$

whereas $\sigma=(2,1)$ corresponds to the reverse ordering of the outer corners and so

$$S_{(2,1),\alpha^j}=\{x\in\mathbf{R}^2_{>0}\mid a_{j+1}\leq x_1<a_j, 0\leq x_2<b_{j+1}\},$$



FIGURE 5.1. The staircase S of M in Example 5.2 and the partitions $\{S_{\alpha^j}\}_j = \{S_{\sigma,\alpha^j}\}_j$ of S corresponding to the permutations $\sigma = (1,2)$ and $\sigma = (2,1)$, respectively.



FIGURE 5.2. The staircase of M in Example 5.3 and the partitions $\{S_{\alpha^j}\}_j$ corresponding to the orderings $\alpha^1, \alpha^2, \alpha^3, \alpha^4, \alpha^5$ and $\alpha^2, \alpha^5, \alpha^3, \alpha^4, \alpha^1$, respectively.

see Figure 5.1. Thus, the partitions just correspond to vertical and horisontal, respectively, slicing of S.

In this case the Scarf complex is a line segment, and it is not very hard to directly compute $d\varphi$, cf. [LW, Section 7]. □

Example 5.3. Let M be the generic monomial ideal $M = (z_1^3, z_1^2 z_2, z_1 z_2^2 z_3^2, z_2^4, z_2^3 z_3, z_3^3)$. The staircase S of M, depicted in Figure 5.2, has five outer corners, $\alpha^1 = (3,1,3)$, $\alpha^2 = (2,4,1)$, $\alpha^3 = (2,3,2)$, $\alpha^4 = (2,2,3)$, and $\alpha^5 = (1,3,3)$. The six different permutations σ of $\{1,2,3\}$ give rise to six different orderings of the α^j: for example $\sigma^1 := (1,2,3)$ and $\sigma^2 := (2,3,1)$ correspond to the orderings $\alpha^1 \geq_\sigma \alpha^2 \geq_\sigma \alpha^3 \geq_\sigma \alpha^4 \geq_\sigma \alpha^5$ and $\alpha^2 \geq_\sigma \alpha^5 \geq_\sigma \alpha^3 \geq_\sigma \alpha^4 \geq_\sigma \alpha^1$, respectively. In the first case S_{σ^1,α^1} is the cuboid $]0,3]\times]0,1]\times]0,3]$, $S_{\sigma^1,\alpha^2} =]0,2]\times]1,4]\times]0,1]$, $S_{\sigma^1,\alpha^3} =]0,2]\times]1,3]\times]1,2]$, $S_{\sigma^1,\alpha^4} =]0,2]\times]1,2]\times]2,3]$, and $S_{\sigma^1,\alpha^5} =]0,1]\times]2,3]\times]2,3]$, see Figure 5.2, where also the S_{σ^2,α^j} are depicted.

□

6. GENERAL (MONOMIAL) IDEALS

The Scarf resolution is an instance of a more general construction of so-called *cellular resolutions* of monomial ideals, introduced by Bayer-Sturmfels [BS]. The Scarf complex is then replaced by a more general oriented polyhedral cell complex X, with vertices corresponding to and labeled by the generators of the monomial ideal M; as above a face γ of X is labeled by the least common multiple m_γ of the vertices. Analogously to the Scarf complex X encodes a graded complex of free A-modules: for $k = 0, \ldots, \dim X + 1$, let E_k be a free A-module of rank equal to the



FIGURE 6.1. The staircase of M in Example 6.1, the partitions S_{α^1} and S_{α^2} corresponding to the orderings α^1, α^2 and α^2, α^1, respectively, and the hull complex of M.

number of $(k-1)$-dimensional faces of X and let $\varphi_k : E_k \to E_{k-1}$ be defined by $\varphi_k : e_\gamma \mapsto \sum_{\delta\subset\gamma} \operatorname{sgn}(\delta,\gamma)\, \frac{m_\gamma}{m_\delta} e_\delta$, where γ and δ are faces of X of dimension $k-1$ and $k-2$, respectively, and where $\operatorname{sgn}(\delta,\gamma) = \pm 1$ comes from the orientation of X. The complex $E_\bullet, \varphi_\bullet$ is exact if X satisfies a certain acyclicity condition see, e.g., [MS, Proposition 4.5], and thus gives a resolution - a so-called *cellular resolution* - of the cokernel of φ_0, which, with the identification $E_{-1} = A$ equals A/M. For more details we refer to [BS] or [MS].

In [BS] was also introduced a certain canonical choice of X. Given $t \in \mathbf{R}$, let $\mathcal{P}_t = \mathcal{P}_t(M)$ be the convex hull in $\mathbf{R}^n$ of $\{(t^{\alpha_1}, \dots, t^{\alpha_n}) \mid z^\alpha \in M\}$. Then $\mathcal{P}_t$ is a unbounded polyhedron in $\mathbf{R}^n$ of dimension n and the face poset (i.e., the set of faces partially ordered by inclusion) of bounded faces of $\mathcal{P}_t$ is independent of t if $t \gg 0$. The *hull complex* of M is the polyhedral cell complex of all bounded faces of $\mathcal{P}_t$ for $t \gg 0$. The corresponding complex $E_\bullet, \varphi_\bullet$ is exact and thus gives a resolution, the *hull resolution*, of A/M. It is in general not minimal, but it has length at most n. If M is generic, however, the Hull complex coincides with the Scarf complex; in particular it is minimal.

In [LW] together with Lärkäng we computed the residue current R associated with the hull resolution, or any cellular resolution where the underlying polyhedral complex X is a polyhedral subdivision of the $(n-1)$-simplex, of an Artinian monomial ideal. Theorem 5.1 in [LW] states that R is of the form (1.9), where the sum is now over all top-dimensional faces (with label α) of X and $\operatorname{sgn}(\alpha)$ comes from the orientation of X.

Note that the definition of $S_{\sigma,\alpha}$ still makes sense when M is a general Artinian monomial ideal. However, in general the $S_{\sigma,\alpha}$ will not be cuboids as the following example shows.

Example 6.1. Let $M = (z_1^2, z_1z_2, z_1z_3, z_2^2, z_3^2)$. Then M is not generic, since there is no generator that strictly divides $\operatorname{lcm}(z_1z_2, z_1z_3) = z_1z_2z_3$.

The staircase S of M is depicted in Figure 6.1. Note that S has two outer corners $\alpha^1 = (2,2,1)$ and $\alpha^2 = (1,1,2)$.

Assume that σ is a permutation of $\{1,2,3\}$ such that $\sigma(1)$ equals 1 or 2. Then the lexicographical order of the outer corners is $\alpha^1 \geq_\sigma \alpha^2$. Otherwise, if $\sigma(1) = 3$, the lexicographical order is reversed. In the first case S_{α^1} is the cuboid $]0,2]\times]0,2]\times]0,1]$, and S_{α^2} is the cuboid $]0,1]\times]0,1]\times]1,2]$, see Figure 6.1. In the second case S_{α^2} is the cuboid $]0,1]\times]0,1]\times]0,2]$, whereas S_{α^1} is the set $]0,2]\times]0,2]\times]0,1]\setminus]0,1]\times]0,1]\times]0,1]$; in particular S_{α^1} is not a cuboid.

In this case, the hull resolution is a minimal resolution of A/M. There are two top-dimensional faces in the hull complex, with vertices $\{z_1^2, z_1z_2, z_1z_3, z_2^2\}$ and $\{z_1z_2, z_1z_3, z_3^2\}$ and thus labels z^{α^1} and z^{α^2}, respectively, see Figure 6.1. A computation yields that if $\sigma(1) = 3$, then the coefficient of $\operatorname{sgn}(\alpha^1)z^{\alpha^1-\mathbf{1}}dze^*_{\alpha^1}$ is $3 = \operatorname{Vol}(S_{\sigma,\alpha^1})$ and the coefficient of $\operatorname{sgn}(\alpha^2)z^{\alpha^2-\mathbf{1}}dze^*_{\alpha^2}$ is $2 = \operatorname{Vol}(S_{\sigma,\alpha^2})$. Otherwise the coefficients are $4 = \operatorname{Vol}(S_{\sigma,\alpha^1})$ and $1 = \operatorname{Vol}(S_{\sigma,\alpha^2})$, respectively. Thus in this case Theorem 1.1 holds. □

Example 6.1 suggests that Theorem 1.1 might hold when (1.2) is the hull resolution of an Artinian monomial and this resolution is minimal. However, we do not know how to prove it in general. The proof in Section 4 does not extend to this situation. For example the staircases R_a constructed in Section 4.1 are not disjoint in general, cf. Claim 4.5. Consider the inner corners $a = (1, 1, 0)$ and $b = (1, 0, 1)$ of the staircase S in Example 6.1. Then

$$R_a \cap R_b = \{x \in \mathbf{R}^3 | x_1 = 1, 1 < x_j \leq 2, j = 2, 3\}.$$

Also, the computation of $d\varphi$ is more involved in this case. Indeed, in general it is not true that the coefficient of $e^*_{\mathcal{I}}$ in (1.8) just consists of one term as in (4.5).

Example 6.2 below shows that Theorem 1.1 does not hold unless the hull resolution is minimal, and also that it does not hold for arbitrary minimal resolutions of monomial ideals. Still in all examples we have computed Corollary 1.2 holds, which motivates the question whether (1.10) holds for all resolutions of minimal length. Note that this question can be asked also for resolutions $E_\bullet, \varphi_\bullet$ of general Artinian ideals $\mathfrak{a}$ in A. One could also ask for a description of the form $d\varphi$ in terms of the staircase (of an initial ideal) of $\mathfrak{a}$.

Example 6.2. Let $M = (z_1^3, z_1^2z_2^2, z_1z_3, z_2^3, z_2z_3, z_3^2)$. Then M is not generic; for example, as in Example 6.1, there is no generator of M that strictly divides $\operatorname{lcm}(z_1z_2, z_1z_3) = z_1z_2z_3$. The staircase of M has three outer corners, $\alpha^1 = (3, 2, 1)$, $\alpha^2 = (2, 3, 1)$, and $\alpha^3 = (1, 1, 2)$.

In this case the hull resolution is not minimal. The hull complex consists of four triangles; one triangle α^j for each outer corner and one extra triangle β with vertices $\{z_1^2z_2^2, z_1z_3, z_2z_3\}$ and thus label $z^\beta = z_1^2z_2^2z_3$, see Figure 6.2. A computation yields that the coefficient of $\operatorname{sgn}(\alpha^3)z^{\alpha^3-\mathbf{1}}dze^*_{\alpha^3}$ in $d_\sigma\varphi$ equals $\operatorname{Vol}(S_{\sigma,\alpha^3})$ for all σ. The coefficient of $\operatorname{sgn}(\alpha^1)z^{\alpha^1-\mathbf{1}}dze^*_{\alpha^1}$ in $d_{(3,1,2)}\varphi$, however, equals 4, whereas $\operatorname{Vol}(S_{(3,1,2),\alpha^1}) = 5$. By symmetry, the coefficient of $\operatorname{sgn}(\alpha^2)z^{\alpha^2-\mathbf{1}}dze^*_{\alpha^2}$ in $d_{(3,2,1)}\varphi$ is 4 and $\operatorname{Vol}(S_{(3,2,1),\alpha^2}) = 5$. Thus Theorem 1.1 does not hold in this case.

However, Corollary 1.2 still holds. This follows using (1.9) , since the coefficient of $\operatorname{sgn}(\beta)z^{\beta-\mathbf{1}}dze^*_\beta$ in $d_\sigma\varphi$ is 1 for $\sigma = (3, 1, 2)$ and $\sigma = (3, 2, 1)$ and 0 otherwise, cf. the beginning of this section.

One can create a minimal cellular resolution from the hull complex, e.g., by removing the edge between $z_1^2z_2^2$ and z_1z_3. The polyhedral cell complex X so obtained has one top-dimensional face for each outer corner α^j in S. The face corresponding to α^1 is the union of the two triangles α^1 and β in the hull complex, see Figure 6.2. It turns out that the coefficient of $\operatorname{sgn}(\alpha^1)z^{\alpha^1-\mathbf{1}}dze^*_{\alpha^1}$ in $d_\sigma\varphi$ is the sum of the coefficients of $\operatorname{sgn}(\alpha^1)z^{\alpha^1-\mathbf{1}}dze^*_{\alpha^1}$ and $\operatorname{sgn}(\beta)z^{\beta-\mathbf{1}}dze^*_\beta$ for each σ, whereas the coefficients of $\operatorname{sgn}(\alpha^2)z^{\alpha^2-\mathbf{1}}dze^*_{\alpha^2}$ and $\operatorname{sgn}(\alpha^3)z^{\alpha^3-\mathbf{1}}dze^*_{\alpha^3}$ are the same as above. In particular, this means that the coefficients of $\operatorname{sgn}(\alpha^j)z^{\alpha^j-\mathbf{1}}dze^*_{\alpha^j}$ equal $\operatorname{Vol}(S_{\sigma,\alpha^j})$ for each α^j



Figure 6.2. The hull complex of the ideal M in Example 6.2 (labels on vertices and 2-faces) (left) and the minimal free resolution (right).

for all σ except $\sigma = (3,2,1)$, in which case the coefficient of $\operatorname{sgn}(\alpha^1)z^{\alpha^1-\mathbf{1}}dze^*_{\alpha^1}$, $\operatorname{sgn}(\alpha^2)z^{\alpha^2-\mathbf{1}}dze^*_{\alpha^2}$, and $\operatorname{sgn}(\alpha^2)z^{\alpha^3-\mathbf{1}}dze^*_{\alpha^3}$ equal $\operatorname{Vol}(S_{\sigma,\alpha^1})+1$ $\operatorname{Vol}(S_{e_{\alpha^2}})-1$, and $\operatorname{Vol}(S_{e_{\alpha^3}})$, respectively. To sum up, Corollary 1.2 holds but Theorem 1.1 fails to hold also in this case.

$\square$

References

[AW] M. Andersson & E. Wulcan: *Residue currents with prescribed annihilator ideals*, Ann. Sci. École Norm. Sup. **40** (2007) no. 6, 985–1007.

[AL-J] B. Angéniol & M. Lejeune-Jalabert: *Calcul différentiel et classes caractéristiques en géométrie algébrique*, Travaux en Cours, **38** Hermann, Paris, 1989.

[BS] D. Bayer & B. Sturmfels: *Cellular resolutions of monomial modules*, J. Reine Angew. Math. **502** (1998) 123–140.

[BPS] D. Bayer & I. Peeva & B. Sturmfels: *Monomial resolutions*, Math. Res. Lett. **5** (1998), no. 1-2, 31–46.

[CH] N. Coleff & M. Herrera: *Les courants résiduels associés à une forme méromorphe*, Lect. Notes in Math. **633**, Berlin-Heidelberg-New York (1978).

[D] Demailly, J.-P.: *Complex and Differential geometry*, available at http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf.

[DP] J.-P. Demailly & M. Passare: *Courants résiduels et classe fondamentale*, Bull. Sci. Math. **119** (1995), no. 1, 85–94.

[E] D. Eisenbud: *The geometry of syzygies. A second course in commutative algebra and algebraic geometry*, Graduate Texts in Mathematics, 229. Springer-Verlag, New York, 2005.

[F] W. Fulton: *Intersection theory. Ergebnisse der Mathematik und ihrer Grenzgebiete*, Springer-Verlag, Berlin, 1984.

[GH] P. Griffiths & J. Harris: *Principles of algebraic geometry*, Pure and Applied Mathematics. Wiley-Interscience, New York, 1978.

[L] R. Lärkäng: *A comparison formula for residue currents*, Preprint, arXiv:1207.1279.

[LW] R. Lärkäng & E. Wulcan: *Computing residue currents of monomial ideals using comparison formulas*, Preprint, arXiv:1205.4648.

[LW2] R. Lärkäng & E. Wulcan: *Residue currents and fundamental cycles*, In preparation.

[L-J] M. Lejeune-Jalabert: *Remarque sur la classe fondamentale d'un cycle*, C. R. Acad. Sci. Paris Sér. I Math. **292** (1981), no. 17, 801–804.

[L-J2] M. Lejeune-Jalabert: *Liaison et résidu*, Algebraic geometry (La Rábida, 1981), 233–240, Lecture Notes in Math., 961, Springer, Berlin, 1982.

[MS] E. Miller & B. Sturmfels: *Combinatorial commutative algebra*, Graduate Texts in Mathematics **227** Springer-Verlag, New York, 2005.

[MSY] E. Miller & B. Sturmfels & K. Yanagawa: *Generic and cogeneric monomial ideals, Symbolic computation in algebra, analysis, and geometry (Berkeley, CA, 1998)*, J. Symbolic Comput. **29** (2000) no 4–5, 691–708.

[T] J. L. Taylor: *Several complex variables with connections to algebraic geometry and Lie groups*, Graduate Studies in Mathematics, **46** American Mathematical Society, Providence, RI, 2002.

Department of Mathematics, Chalmers University of Technology and the University of Gothenburg, S-412 96 Gothenburg, SWEDEN
E-mail address: `wulcan@chalmers.se`